


GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:



GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco media and advertising, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is self-administered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In Turkey, GYTS was conducted in 2017 by the Turkish Institute of Public Health, under the coordination of the Ministry of Health. The overall response rate was 82.8%. A total of 122,040 eligible students in grades secondary 7, secondary 8, preparatory, high school 1, and high school 2 completed the survey, of which 84,007 were aged 13-15 years. Data are reported for students aged 13-15 years.

GYTS Highlights

TOBACCO USE

- 17.9% of students, 23.2% of boys, and 12.1% of girls currently used any tobacco products.
- 17.9% of students, 23.2% of boys, and 12.1% of girls currently smoked tobacco.
- 7.7% of students, 9.9% of boys, and 5.3% of girls currently smoked cigarettes.

CESSATION

- Almost 6 in 10 (56.4%) current tobacco smokers tried to stop smoking in the past 12 months.
- More than 5 in 10 (53.2%) current tobacco smokers wanted to stop smoking now.

SECONDHAND SMOKE

- 46.1% of students were exposed to tobacco smoke at home.
- 51.8% of students were exposed to tobacco smoke inside enclosed public places.

ACCESS & AVAILABILITY

- 61.6% of current cigarette smokers bought cigarettes from a store, shop, or kiosk (bufe).
- Among current cigarette smokers who tried to buy cigarettes, 73.3% were not prevented from buying them because of their age.

MEDIA

- More than 6 in 10 (61.1%) students noticed anti-tobacco messages in the media.
- Almost 3 in 10 (26.8%) students noticed tobacco advertisements or promotions when visiting points of sale.
- Almost 1 in 10 (8.3%) students had something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 79.5% of students definitely thought other people's tobacco smoking is harmful to them.
- 91.1% of students favored prohibiting smoking inside enclosed public places.
- 74.5% of students favored prohibiting smoking at outdoor public places.

TOBACCO USE


SMOKED TOBACCO	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers ¹	17.9	23.2	12.1*
Current cigarette smokers ²	7.7	9.9	5.3*
Frequent cigarette smokers ³	2.3	3.7	0.9*
Current smokers of other tobacco ⁴	12.8	17.3	8.0*
Ever tobacco smokers ⁵	40.2	46.8	33.1*
Ever cigarette smokers ⁶	28.0	32.9	23.0*
Ever smokers of other tobacco ⁷	23.2	29.5	16.5*

ANY TOBACCO USE (smoked tobacco only)⁸

Current tobacco users ¹	17.9	23.2	12.1*
Ever tobacco users ⁵	40.2	46.8	33.1*

SUSCEPTIBILITY

Never tobacco users susceptible to tobacco use in the future ⁹	9.9	10.6	9.2
Never tobacco smokers who thought they might enjoy smoking a cigarette ¹⁰	12.9	14.9	11.1*



CESSATION

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers who tried to stop smoking in the past 12 months	56.4	61.1	46.8*
Current tobacco smokers who wanted to stop smoking now	53.2	57.4	43.7*
Current tobacco smokers who thought they would be able to stop smoking if they wanted to	67.3	62.7	77.2*
Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking	13.6	15.7	9.3*

SECONDHAND SMOKE

	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home ¹¹	46.1	44.1	48.3*
Exposure to tobacco smoke inside any enclosed public place ¹¹	51.8	50.9	52.6*
Exposure to tobacco smoke at any outdoor public place ¹¹	54.2	51.3	57.2*
Students who saw anyone smoking inside the school building or outside on school property ¹²	59.3	59.5	59.2

ACCESS & AVAILABILITY

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current cigarette smokers who bought cigarettes from a store, shop, or kiosk (bufe) ¹³	61.6	63.8	58.8
Current cigarette smokers who were not prevented from buying cigarettes because of their age ¹⁴	73.3	71.5	76.7
Current cigarette smokers who bought cigarettes as individual sticks ¹⁵	29.4	29.7	28.4

MEDIA

TOBACCO ADVERTISING	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who noticed tobacco advertisements or promotions at points of sale ¹⁶	26.8	28.9	24.5*
Students who saw anyone using tobacco on television, videos, or movies ¹⁷	66.5	66.1	67.0
Students who were ever offered a free tobacco product from a tobacco company representative	8.6	12.8	4.4*
Students who had something with a tobacco brand logo on it	8.3	11.7	5.0*

ANTI-TOBACCO ADVERTISING

Students who noticed anti-tobacco messages in the media ¹²	61.1	58.5	63.8*
Students who noticed anti-tobacco messages at sporting or community events ¹⁸	35.2	37.1	32.9*
Current tobacco smokers who thought about quitting because of a warning label ¹⁹	22.9	24.0	20.5*
Students who were taught in school about the dangers of tobacco use in the past 12 months	56.0	54.7	57.3*

KNOWLEDGE & ATTITUDES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	36.9	33.6	40.3*
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	26.2	31.0	21.3*
Students who definitely thought other people's tobacco smoking is harmful to them	79.5	75.1	84.2*
Students who favored prohibiting smoking inside enclosed public places	91.1	88.1	94.2*
Students who favored prohibiting smoking at outdoor public places	74.5	72.8	76.3*

¹ Smoked cigarettes or other type of tobacco anytime during the past 30 days. ² Smoked cigarettes anytime during the past 30 days. ³ Smoked cigarettes on 20 or more days of the past 30 days. ⁴ Smoked tobacco other than cigarettes anytime during the past 30 days. ⁵ Ever smoked cigarettes or other type of tobacco, even one or two puffs. ⁶ Ever smoked cigarettes, even one or two puffs. ⁷ Ever smoked tobacco other than cigarettes, even one or two puffs. ⁸ Turkey did not collect data on smokeless tobacco use in this survey. ⁹ Susceptible to future tobacco use includes those who answered "definitely yes", "probably yes", or "probably not" to using tobacco if one of their best friends offered it to them, or "definitely yes", "probably yes", or "probably not" to using tobacco during the next 12 months. ¹⁰ Those who answered "Agree" or "Strongly Agree" to the statement: "I think I might enjoy smoking a cigarette". ¹¹ During the past 7 days. ¹² During the past 30 days. ¹³ Outlet from which current cigarette smokers bought cigarettes the last time they smoked cigarettes in the past 30 days. ¹⁴ Among those who tried to buy cigarettes during the past 30 days. ¹⁵ Based on the last purchase, among those who bought cigarettes during the past 30 days. ¹⁶ Among those who visited a point of sale in the past 30 days. ¹⁷ Among those who watched television, videos, or movies in the past 30 days. ¹⁸ Among those who attended sporting or community events in the past 30 days. ¹⁹ Among those who noticed warning labels on cigarette packages in the past 30 days.

NOTE: Data are weighted to be nationally representative of all students aged 13-15 years who are enrolled in school. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.

*Gender comparisons are significant at p < 0.05.

†The nonlinear increase presented in the graph is significant at p < 0.05 from 2003-2017.